

Value-added reporting

X. Bossuyt

COMMUNICATING DIAGNOSTIC ACCURACY

Communicating diagnostic accuracy

Question 1

- Sensitivity: 95%
- Specificity: 90%
- Pre-test probability: 2.5%
- Post-test probability ???

- 10%
- 20%
- 50%
- 80%
- 90%

Communicating diagnostic accuracy

Question 1

- Sensitivity: 95%
- Specificity: 90%
- Pre-test probability: 2.5%
- Post-test probability ???

- 10%
- 20%
- 50%
- 80%
- 90%

Communicating diagnostic accuracy

Question 1

- Sensitivity: 95%
- Specificity: 90%
- Pre-test probability: 2.5%
- Post-test probability ???

- 10%
- **20% (9% of respondents)**
- 50%
- 80%
- **90%**

Communicating diagnostic accuracy

Question 2

- LR 10

A pos test result occurs 10 times more frequently in patients with the disease compared to patients without the disease

- Pre-test probability: 2.5%

- 10%
- 20%
- 50%
- 80%
- 90%

Communicating diagnostic accuracy

Question 2

- LR 10


A pos test result occurs 10 times more frequently in patients with the disease compared to patients without the disease

- Pre-test probability: 2.5%

- 10%
- **20% (35% of respondents)**
- 50%
- 80%
- 90%

Communicating diagnostic accuracy

- **Question 3**




RHEUMATOID ARTHRITIS


Anti-CCP


298 patients


- RA patients (n=85)
 - 45 early RA
 - 40 established RA
- Diseased controls (n=165), including:
 - Psoriatic arthritis
 - Connective tissue disease
 - Organ-specific autoimmune disease
- Consecutive patients (n=48)
 - 16 RA

- Anti-CCP: Phadia (7Units/mL)
- RF: nephelometry, Immage, Beckman-Coulter (20 IU/mL)









LR

1

no clinical value

>10 <0.1


clinically important difference


5 – 10 0.1 – 0.2


modest, but substantial difference


2 – 5 0.2 – 0.5


small difference that may be relevant

























Female, 50 years old
Recent onset undifferentiated arthritis
Intermittent asymmetric tender and swollen small joints of the hands (n=5)
CRP: 10 mg/L



Female, 50 years old

Symmetric arthritis of upper and lower extremities

Tender and swollen small joints (n=12)

CRP: 70 mg/L

Conclusions

- We showed that a graphical representation is the best way to convey information on post-test probabilities.
- We illustrated how likelihood ratios depend on antibody level.
- Such knowledge may help with the interpretation of a specific test result.
- Clinical laboratories might consider to provide likelihoods ratios or probability data for autoantibody test result intervals.

Acknowledgments

- Department of laboratory medicine
 - P. Vermeersch
 - D. Coenen
- Internal medicine, rheumatology
 - P. Verschueren
 - R. Westhovens

References

- Vermeersch P and Bossuyt X.
Comparative analysis of different approaches to report diagnostic accuracy.
Arch Intern Med. 2010;170:734-5.
- Bossuyt X et al.
Likelihood ratios as a function of antibody concentration for anti-cyclic citrullinated peptide antibodies and rheumatoid factor.
Ann Rheum Dis. 2009;68:287-9.